



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2149052 C 2003/03/25

(11)(21) **2 149 052**

(12) **BREVET CANADIEN  
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1993/11/17  
(87) Date publication PCT/PCT Publication Date: 1994/06/09  
(45) Date de délivrance/Issue Date: 2003/03/25  
(85) Entrée phase nationale/National Entry: 1995/05/09  
(86) N° demande PCT/PCT Application No.: US 1993/011172  
(87) N° publication PCT/PCT Publication No.: 1994/012200  
(30) Priorité/Priority: 1992/12/02 (984,399) US

(51) Cl.Int.<sup>6</sup>/Int.Cl.<sup>6</sup> A61K 31/66, A61K 47/38, A61K 47/26,  
A61K 9/20

(72) Inventeurs/Inventors:  
BECHARD, SIMON R., CA;  
KRAMER, KENNETH A., US;  
KATDARE, ASHOK V., US

(73) Propriétaire/Owner:  
MERCK & CO., INC., US

(74) Agent: OGILVY RENAULT

(54) Titre : FORMULATIONS SOUS FORME DE MELANGE SEC POUR ACIDES BISPHOSPHONIQUES

(54) Title: DRY MIX FORMULATION FOR BISPHOSPHONIC ACIDS

(57) Abrégé/Abstract:

Pharmaceutical compositions of bisphosphonic acids, and salts thereof, are prepared by direct compression/dry mix tablet formulation. These pharmaceutical compositions are useful in the treatment of disturbances involving calcium or phosphate metabolism, in particular, the treatment and prevention of diseases involving bone resorption, especially osteoporosis, Paget's disease, malignant hypercalcemia, and metastatic bone disease.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : A61K 37/00, 31/66, 31/045, 47/00, 9/36 2149052		A1	(11) International Publication Number: WO 94/12200 (43) International Publication Date: 9 June 1994 (09.06.94)
(21) International Application Number: PCT/US93/11172 (22) International Filing Date: 17 November 1993 (17.11.93) (30) Priority Data: 984,399 2 December 1992 (02.12.92) US (60) Parent Application or Grant (63) Related by Continuation US 984,399 (CIP) Filed on 2 December 1992 (02.12.92) (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BECHARD, Simon, R. [CA/CA]; 1755 Marchand, Laval, Quebec H7G 4V7 (CA). KRAMER, Kenneth, A. [US/US]; 426 Sterners Road, Green Lane, PA 18054 (US). KATDARE, Ashok. V. [US/US]; 801 East Fomance Street, Norristown, PA 19401 (US). (74) Agent: WINOKUR, Melvin; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published With international search report.
(54) Title: DRY MIX FORMULATION FOR BISPHOSPHONIC ACIDS			
(57) Abstract  Pharmaceutical compositions of bisphosphonic acids, and salts thereof, are prepared by direct compression/dry mix tablet formulation. These pharmaceutical compositions are useful in the treatment of disturbances involving calcium or phosphate metabolism, in particular, the treatment and prevention of diseases involving bone resorption, especially osteoporosis, Paget's disease, malignant hypercalcemia, and metastatic bone disease.			

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A process for the preparation of a tablet containing an active ingredient selected from the group consisting of:  
 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
 N-methyl-4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
 4-(N,N-dimethylamino)-1-hydroxybutylidene-1,1-bisphosphonic acid;  
 3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid;  
 3-(N,N-dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid;  
 1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid;  
 1-hydroxy-2-[3-pyridyl]ethylidene-1,1-bisphosphonic acid;  
 and  
 4-(hydroxymethylene-1,1-bisphosphonic acid)piperidine;  
 or a pharmaceutically acceptable salt thereof;  
 which process comprises:  
 forming a mixture by mixing the active ingredient with:  
 a diluent selected from: anhydrous lactose and hydrous fast flow lactose,  
 a dry binder, and  
 a disintegrant,  
 lubricating the mixture with a lubricant; and  
 compressing the resultant lubricated mixture into a desired tablet form.
2. The process of claim 1, wherein the active ingredient is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid.
3. The process of claim 1, wherein the active ingredient is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium salt trihydrate.
4. The process of claim 1, 2 or 3, wherein the dry binder is microcrystalline cellulose.

5. The process of claim 1, 2, 3 or 4, wherein the disintegrant is selected from the group consisting of modified starch, modified cellulose polymer and croscarmallose sodium, or a combination thereof.
6. The process of claim 1, 2, 3, 4 or 5, wherein the disintegrant is croscarmallose sodium.
7. The process of claim 1, 2, 3, 4, 5 or 6, wherein the lubricant is magnesium stearate.
8. The process of claim 1, 2, 3, 4, 5, 6 or 7, wherein said mixture further includes one or more additional ingredients selected from the group consisting of: compression aids, flavors, flavor enhancers, sweeteners and preservatives.
9. The process of claim 1, 2, 3, 4, 5, 6, 7 or 8, wherein said diluent is anhydrous lactose.
10. The process of claim 1, 2, 3, 4, 5, 6, 7 or 8, wherein said diluent is hydrous fast flow lactose.
11. A pharmaceutical composition comprising by weight, about 0.5 to 40% by weight of an active ingredient selected from the group consisting of:
  - 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;
  - N-methyl-4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;
  - 4-(N,N-dimethylamino)-1-hydroxybutylidene-1,1-bisphosphonic acid;
  - 3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid;
  - 3-(N,N-dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid;
  - 1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid;
  - 1-hydroxy-2-[3-pyridyl]ethylidene-1,1-bisphosphonic acid;
  - and 4-(hydroxymethylene-1,1-bisphosphonic acid)piperidine;

or a pharmaceutically acceptable salt thereof;  
and from about 60 to 99.5% by weight of excipients consisting essentially of:  
anhydrous lactose or hydrous fast flow lactose; microcrystalline cellulose;  
croscarmallose sodium; and magnesium stearate.

12. A composition of claim 11, containing said anhydrous lactose.

13. A composition of claim 11, containing said hydrous fast flow lactose.

14. A pharmaceutical composition comprising by weight, about 0.5 to 40% by weight of an active ingredient selected from the group consisting of:

4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
N-methyl-4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
4-(N,N-dimethylamino)-1-hydroxybutylidene-1,1-bisphosphonic acid;

3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid;  
3-(N,N-dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid;

1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid;

1-hydroxy-2-[3-pyridyl]ethylidene-1,1-bisphosphonic acid;  
and 4-(hydroxymethylene-1,1-bisphosphonic acid)piperidine;

or a pharmaceutically acceptable salt thereof;

about 10 to 80% by weight of anhydrous lactose or hydrous fast flow lactose;

about 5 to 50% by weight of microcrystalline cellulose;

about 0.5 to 10% by weight of croscarmallose sodium; and

about 0.1 to 5% by weight of magnesium stearate.

15. The pharmaceutical composition of Claim 14, comprising about 0.5 to 25% by weight of the active ingredient, about 30 to 70% by weight of anhydrous lactose or hydrous fast flow lactose, about 30 to 50% by weight of microcrystalline cellulose; about 0.5 to 5% by weight of croscarmallose sodium and about 0.1 to 2% by weight of magnesium stearate.

16. The pharmaceutical composition of claim 14 or 15, containing about 30 to 70% by weight of said anhydrous lactose.
17. The pharmaceutical composition of claim 14 or 15, containing about 30 to 70% by weight of said hydrous fast flow lactose.
18. The pharmaceutical composition of claim 14, 15, 16 or 17, wherein the active ingredient is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid.
19. The pharmaceutical composition of claim 14, 15, 16 or 17, wherein the active ingredient is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium salt trihydrate.
20. The pharmaceutical composition of claim 14 or 15, comprising about 1 to 25% by weight of 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium salt trihydrate as said active ingredient, about 40 to 60% by weight of anhydrous lactose; about 35 to 45% by weight of microcrystalline cellulose; about 0.5 to 2% by weight of croscarmallose sodium and about 0.1 to 1% by weight of magnesium stearate.
21. A tablet prepared from the pharmaceutical composition of claim 14, 15, 16, 17, 18, 19 or 20.
22. A pharmaceutical composition comprising by weight, about 0.5 to 40% weight of an active ingredient selected from the group consisting of:  
4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
N-methyl-4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
4-(N,N-dimethylamino)-1-hydroxybutylidene-1,1-bisphosphonic acid;  
3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid;  
3-(N,N-dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid;  
1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid;

1-hydroxy-2-[3-pyridyl]ethylidene-1,1-bisphosphonic acid;  
and

4-(hydroxymethylene-1,1-bisphosphonic acid)piperidine;

or a pharmaceutically acceptable salt thereof; and from about 60 to 99.5% by weight of excipients comprising:

a diluent selected from: anhydrous lactose and hydrous fast flow lactose; a dry binder, a disintegrant and a lubricant.

23. A pharmaceutical composition comprising from about 0.5 to about 40% by weight of a bisphosphonic acid or a pharmaceutically acceptable salt thereof and from about 60% to 99.5% by weight of excipients, said excipients comprising a diluent selected from: anhydrous lactose and hydrous fast flow lactose; a dry binder, a disintegrant and a lubricant.

24. A pharmaceutical composition according to claim 23, wherein said bisphosphonic acid or pharmaceutically acceptable salt thereof is a nitrogen containing bisphosphonic acid or a pharmaceutically acceptable salt thereof.

25. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.

26. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium salt trihydrate.

27. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is N-methyl-4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.

28. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 4-(N,N-dimethylamino)-1-hydroxybutylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.
29. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.
30. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 3-(N,N-dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.
31. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.
32. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 1-hydroxy-2-[3-pyridyl]ethylidene-1,1-bisphosphonic acid or a pharmaceutically acceptable salt thereof.
33. A pharmaceutical composition according to claim 24, wherein said nitrogen containing bisphosphonic acid or pharmaceutically acceptable salt thereof is 4-(hydroxymethylene-1,1-bisphosphonic acid) piperidine or a pharmaceutically acceptable salt thereof.
34. A pharmaceutical composition according to any one of claims 22 to 33, wherein said composition is coated.
35. A pharmaceutical composition according to claim 34 wherein said composition is coated with a substance selected from the group consisting

of hydroxypropylmethylcellulose, hydroxypropylcellulose, titanium oxide, talc, sweeteners, and colorants.

36. A pharmaceutical composition according to any one of claims 22 to 35 wherein said diluent is anhydrous lactose.

37. A pharmaceutical composition according to any one of claims 22 to 36 wherein said composition is a tablet.

38. A process for the preparation of a tablet containing as active ingredient a basic nitrogen containing bisphosphonate: which process comprises:

forming a mixture by mixing the active ingredient with:  
a diluent selected from: anhydrous lactose and hydrous fast flow lactose,  
a dry binder, and  
a disintegrant,  
lubricating the mixture with a lubricant; and  
compressing the resultant lubricated mixture into a desired tablet form.

39. A process according to claim 38, wherein said mixture further includes one or more additional ingredients selected from the group consisting of: compression aids, flavors, flavor enhancers, sweeteners and preservatives.

40. A process according to claim 38 or 39, wherein said diluent is said anhydrous lactose.

41. A process according to claim 38 or 39, wherein said diluent is said hydrous fast flow lactose.

42. A tablet prepared by the process of claim 1, 2, 3, 4, 5, 6, 7, 8 or 9.

2149052

- 1 -

TITLE OF THE INVENTION

DRY MIX FORMULATION FOR BISPHOSPHONIC ACIDS

BACKGROUND OF THE INVENTION

5           The pharmaceutical industry employs various methods for compounding pharmaceutical agents in tablet formulations. In particular, wet granulation is one of the most prevalent methods.

          A variety of bisphosphonic acids have been disclosed as being useful in the treatment and prevention of diseases involving bone  
10       resorption. Representative examples may be found in the following: U.S. Patent No. 3,962,432; U.S. Patent No. 4,054,598; U.S. Patent No. 4,267,108; U.S. Patent No. 4,327,039; U.S. Patent No. 4,621,077; U.S. Patent No. 4,624,947; U.S. Patent No. 4,746,654; U.S. Patent No. 4,922,077; and EPO Patent Pub. No. 0,252,504. Standard methods for  
15       tablet formulation of bisphosphonic acids, however, suffer serious difficulties.

          In particular, bisphosphonic acids which bear a basic nitrogen-containing functionality may interact with the lactose of  
20       standard formulations resulting in discoloration, instability and potency loss. This degradation of the active ingredient is particularly pronounced in the presence of water and/or elevated temperature. It is speculated that this incompatibility is specifically due to the Maillard (or "browning") reaction in which the free amino group of the  
25       bisphosphonic acid reacts with the "glycosidic" hydroxyl group of a sugar (such as lactose) ultimately resulting in the formation of brown pigmented degradates. Although this problem may be avoided by the elimination of lactose, the use of lactose as an inert diluent is generally desirable.

30       The present invention solves this problem by providing a tablet formulation and process therefor that avoids such interaction between the bisphosphonic acid and the lactose in the formulation. In addition, the present invention also provides a processing advantage since it requires only blending of the ingredients without granulation or addition of water prior to compression.

- 2 -

DESCRIPTION OF THE INVENTION

5 The present invention is directed in a first embodiment to a process for the preparation of pharmaceutical compositions of bisphosphonic acids by direct compression (dry mix) tablet formulation. This process employs a blend of a bisphosphonic acid and minimal amounts of other processing aids with no water added. The tablet formulation is prepared by mixing the formulation ingredients with no hydration (i.e. no additional water is added to the mixture) prior to  
10 direct compression.

More specifically, this embodiment of the present invention concerns a process for the preparation of a tablet containing a bisphosphonic acid as an active ingredient which process comprises:  
15 forming a mixture by mixing the active ingredient with:  
a diluent,  
a dry binder,  
a disintegrant,  
and optionally one or more additional ingredients selected from the group consisting of: compression aids, flavors,  
20 flavor enhancers, sweeteners and preservatives;  
lubricating the mixture with a lubricant; and  
compressing the resultant lubricated mixture into a desired tablet form.

25 The disclosed process may be used to prepare solid dosage forms, particularly tablets, for medicinal administration.

Preferred diluents include lactose. In particular, anhydrous lactose is preferred from the flow processing point of view, although  
30 hydrous fast flow lactose may also be employed.

A preferred dry binder is cellulose. In particular, microcrystalline cellulose is preferred. Microcrystalline cellulose is available commercially under the trade mark "Avicel" from FMC Corporation.

- 3 -

The disintegrant is suitably selected from the group consisting of modified starch, modified cellulose polymer and croscarmallose sodium, or a combination thereof and may be one of several modified starches or modified cellulose polymers, in particular, croscarmallose sodium is preferred. Croscarmallose sodium NF Type A (NF refers to the U.S. National Formulary) is commercially available under the trade mark "Ac-di-sol".

Preferred lubricants include magnesium stearate.

Examples of the bisphosphonic acids which may be employed as active ingredients in the instant invention include:

4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
 N-methyl-4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;  
 4-(N,N-dimethylamino)-1-hydroxybutylidene-1,1-bisphosphonic acid;  
 3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid;  
 3-(N,N-dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid;  
 1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid;  
 1-hydroxy-2-[3-pyridyl]ethylidene-1,1-bisphosphonic acid; and  
 4-(hydroxymethylene-1,1-bisphosphonic acid)piperidine;  
 or a pharmaceutically acceptable salt thereof.

Methods for the preparation of bisphosphonic acids may be found in, e.g. U.S. Patent No. 3,962,432; U.S. Patent No. 4,054,598; U.S. Patent No. 4,267,108; U.S. Patent No. 4,327,039; U.S. Patent No. 4,407,761; U.S. Patent No. 4,621,077; U.S. Patent No. 4,624,947; U.S. Patent No. 4,746,654; U.S. Patent No. 4,922,077 and EPO Patent Pub. No. 0,252,504. In particular, methods for the preparation of 4-amino-1-hydroxy-butylidene-1,1-bisphosphonic acid and 4-amino-1-hydroxy-butylidene-1,1-bisphosphonic acid monosodium salt trihydrate may be found in U.S. Patent No. 4,407,761 and U.S. Patent 4,922,077, respectively.

The pharmaceutically acceptable salts of bisphosphonic acids may also be employed in the instant invention. Examples of base

2149052

- 4 -

5 salts of bisphosphonic acids include ammonium salts, alkali metal salts such as potassium and sodium (including mono-, di- and tri-sodium) salts (which are preferred), alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D- glucamine, and salts with amino acids such as arginine, lysine, and so forth. The non-toxic, physiologically acceptable salts are preferred. The salts may be prepared by methods known in the art, such as in U.S. Patent No. 4,922,077.

10 In the present invention it is preferred that the bisphosphonic acid is 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid. It is even more preferred that the bisphosphonic acid is a sodium salt of 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid, in particular, 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium salt trihydrate.

15 Another embodiment of the present invention is a direct compression pharmaceutical composition, such as a tablet, comprising a bisphosphonic acid, which is prepared by the disclosed process. In general, these pharmaceutical compositions comprise by weight, about 0.5 to 40% by weight of a bisphosphonic acid as an active ingredient; and from about 60 to 99.5% by weight of processing aids with no water  
20 added. More specifically, the processing aids are a diluent, a dry binder, a disintegrant and a lubricant. Preferred processing aids include: anhydrous lactose or hydrous fast flow lactose; microcrystalline cellulose; croscarmallose sodium; and magnesium  
25 stearate.

Preferred pharmaceutical compositions comprise about 0.5 to 40% by weight of a bisphosphonic acid as an active ingredient; about 10 to 80% by weight of anhydrous lactose or hydrous fast flow lactose; about 5 to 50% by weight of microcrystalline cellulose; about 0.5 to  
30 10% by weight of croscarmallose sodium; and about 0.1 to 5% by weight of magnesium stearate.

The preferred pharmaceutical compositions are generally in the form of tablets. The tablets may be, for example, from 50 mg to

2149052

- 5 -

1.0 g in net weight, more preferably 100 to 500 mg net weight, and even more preferably 200 to 300 mg net weight.

More preferred pharmaceutical compositions in accordance with the present invention comprise: about 0.5 to 25% by weight of a  
5 bisphosphonic acid selected from 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid and 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium salt trihydrate; about 30 to 70% by weight of anhydrous lactose or hydrous fast flow lactose; about 30 to 50% by  
10 weight of microcrystalline cellulose; about 0.5 to 5% by weight of croscarmallose sodium; and about 0.1 to 2% by weight of magnesium stearate.

Especially preferred pharmaceutical compositions comprise about 1 to 25% of the active ingredient, about 40 to 60% by weight of  
15 anhydrous lactose; about 35 to 45% by weight of microcrystalline cellulose; about 0.5 to 2% by weight of croscarmallose sodium; and about 0.1 to 1% by weight of magnesium stearate. Preferred pharmaceutical compositions as envisioned for commercial development are as follows.

Tablets of 2.5 mg potency free acid:  
20 about 1.63% by weight of 4-amino-1-hydroxy-butylidene-1,1-bisphosphonic acid monosodium salt trihydrate; about 56.87% by weight of anhydrous lactose; about 40% by weight of microcrystalline cellulose; about 1% by weight of croscarmallose sodium; and about  
25 0.5% by weight of magnesium stearate.

Tablets of 5 mg potency free acid:  
about 3.25% by weight of 4-amino-1-hydroxy-butylidene-1,1-bisphosphonic acid monosodium salt trihydrate; about 55.25% by  
weight of anhydrous lactose; about 40% by weight of microcrystalline  
30 cellulose; about 1% by weight of croscarmallose sodium; and about 0.5% by weight of magnesium stearate.

Tablets of 25 mg potency free acid:  
about 16.4% by weight of 4-amino-1-hydroxy-butylidene-1,1-bisphosphonic acid monosodium salt trihydrate; about 42.1% by  
weight of anhydrous lactose; about 40% by weight of microcrystalline

2149052

- 6 -

cellulose; about 1% by weight of croscarmallose sodium; and about 0.5% by weight of magnesium stearate.

Tablets of 50 mg potency free acid:

5       about 21.8% by weight of 4-amino-1-hydroxy-butylidene-1,1-bisphosphonic acid monosodium salt trihydrate; about 36.7% by weight of anhydrous lactose; about 40% by weight of microcrystalline cellulose; about 1% by weight of croscarmallose sodium; and about 0.5% by weight of magnesium stearate.

10       The pharmaceutical tablet compositions of the present invention may also contain one or more additional formulation ingredients may be selected from a wide variety of excipients known in the pharmaceutical formulation art. According to the desired properties of the tablet, any number of ingredients may be selected, alone or in  
15       combination, based upon their known uses in preparing tablet compositions. Such ingredients include, but are not limited to, diluents, compression aids, disintegrants, lubricants, binders, flavors, flavor enhancers, sweetener and preservatives.

20       The term "tablet" as used herein is intended to encompass compressed pharmaceutical dosage formulations of all shapes and sizes, whether coated or uncoated. Substances which may be used for coating include hydroxypropylmethylcellulose, hydroxypropylcellulose, titanium oxide, talc, sweeteners, and colorants.

25       The pharmaceutical compositions of the present invention are useful in the therapeutic or prophylactic treatment of disorders in calcium or phosphate metabolism and associated diseases. These diseases can be divided into two categories:

- 30       1.     Abnormal (ectopic) depositions of calcium salts, mostly calcium phosphate, pathological hardening of tissues and bone malformations.
2.     Conditions which can benefit from a reduction in bone resorption. A reduction in bone resorption should improve the balance between resorption and formation, reduce bone loss or result in bone augmentation. A reduction in bone resorption can alleviate the

- 7 -

pain associated with osteolytic lesions and reduce the incidence and/or growth of those lesions.

5        These diseases include: osteoporosis (including estrogen deficiency, immobilization, glucocorticoid induced and senile), osteodystrophy, Paget's disease, myositis ossificans, Bechterew's disease, malignant hypercalcemia, metastatic bone disease, periodontal disease, cholelithiasis, nephrolithiasis, urolithiasis, urinary calculus, hardening of the arteries (sclerosis), arthritis, bursitis, neuritis and tetany.

10       Increased bone resorption can be accompanied by pathologically high calcium and phosphate concentrations in the plasma, which would be alleviated by use of the instant pharmaceutical compositions.

15       The following examples are given for the purpose of illustrating the present invention and shall not be construed as being limitations on the scope or spirit of the invention.

20

25

30

2149052

- 8 -

EXAMPLE 1

Procedure for Manufacturing 5 mg Potency Tablets of 4-Amino-1-hydroxybutylidene-1,1-bisphosphonic acid

5

<u>Ingredients</u>		<u>Per Tablet</u>	<u>Per 4,000 Tablets</u>
10	Active ingredient (monosodium salt trihydrate)	6.55 mg	26.2 g
	Anhydrous Lactose, NF	110.45 mg	441.8 g
15	Microcrystalline Cellulose NF	80.0 mg	320.0 g
	Magnesium Stearate Impalpable Powder NF	1.00 mg	4.0 g
20	Croscarmellose Sodium NF Type A	2.00 mg	8.0 g

25 The active ingredient (equivalent to 5 mg anhydrous free acid per tablet) was premixed with 1/3 of the microcrystalline cellulose NF and 1/2 of the anhydrous lactose NF in a ribbon blender for 5 minutes at 20 RPM. To the premix was added the remaining 2/3 of the microcrystalline cellulose NF and the remaining 1/2 of the anhydrous lactose NF. This was blended for 10 minutes at 20 RPM.

30 Crosscarmellose sodium was added to the blended powders and mixed for 5 minutes at 20 RPM. Finally the magnesium stearate was added to the mixture by passing through a 90 mesh screen and blended for an additional 5 minutes at 20 RPM. The lubricated mixture was compressed to provide tablets of 5 mg active ingredient.

- 9 -

EXAMPLE 2

Procedure for Manufacturing 2.5 mg Potency Tablets of 4-Amino-1-  
hydroxybutylidene-1,1-bisphosphonic acid

5

<u>Ingredients</u>	<u>Per Tablet</u>
Active ingredient (monosodium salt trihydrate)	3.26 mg
Anhydrous Lactose, NF	113.74 mg
Microcrystalline Cellulose NF	80.0 mg
Magnesium Stearate Impalpable Powder NF	1.00 mg
Croscarmellose Sodium NF Type A	2.00 mg

10

15

20

Tablets were prepared using the procedure of Example 1.

25

30

2149052

- 10 -

EXAMPLE 3

Procedure for Manufacturing 10.0 mg Potency Tablets of 4-Amino-1-  
hydroxybutylidene-1,1-bisphosphonic acid

5

	<u>Ingredients</u>	<u>Per Tablet</u>
10	Active ingredient (monosodium salt trihydrate)	13.05 mg
	Anhydrous Lactose, NF	103.95 mg
15	Microcrystalline Cellulose NF	80.0 mg
	Magnesium Stearate Impalpable Powder NF	1.00 mg
20	Croscarmellose Sodium NF Type A	2.00 mg

Tablets were prepared using the procedure of Example 1.

25

30

- 11 -

EXAMPLE 4

Procedure for Manufacturing 40.0 mg Potency Tablets of 4-Amino-1-  
hydroxybutylidene-1,1-bisphosphonic acid

5

<u>Ingredients</u>	<u>Per Tablet</u>
Active ingredient (monosodium salt trihydrate)	51.21 mg
Anhydrous Lactose, NF	64.79 mg
Microcrystalline Cellulose NF	80.0 mg
Magnesium Stearate Impalpable Powder NF	1.00 mg
Croscarmellose Sodium NF Type A	2.00 mg

10

15

20

Tablets were prepared using the procedure of Example 1.

25

30

2149052

- 12 -

EXAMPLE 5

Procedure for Manufacturing 25 mg Potency Tablets of 4-Amino-1-hydroxybutylidene-1,1-bisphosphonic acid

5

Ingredients	Per Tablet	Per 4,000
		Tablets
Active ingredient (monosodium salt trihydrate)	32.75 mg	131.0 g
Anhydrous Lactose, NF	84.25 mg	337.0 g
Microcrystalline Cellulose NF	80.0 mg	320.0 g
Magnesium Stearate Impalpable Powder NF	1.00 mg	4.0 g
Croscarmellose Sodium NF Type A	2.00 mg	8.0 g

10

15

20

Tablets were prepared using the procedure of Example 1.

25

30

- 13 -

EXAMPLE 6

Procedure for Manufacturing 50 mg Potency Tablets of 4-Amino-1-hydroxybutylidene-1,1-bisphosphonic acid

5	<u>Ingredients</u>	<u>Per Tablet</u>	<u>Per 2,500 Tablets</u>
	Active ingredient (monosodium salt trihydrate)	65.5 mg	163.75 g
10	Anhydrous Lactose, NF	110.0 mg	275.0 g
	Microcrystalline Cellulose NF	120.0 mg	300.0 g
15	Magnesium Stearate Impalpable Powder NF	1.5 mg	3.75 g
20	Croscarmellose Sodium NF Type A	3.0 mg	7.5 g

Tablets were prepared using the procedure of Example 1.

EXAMPLE 7

25

Stability Studies

Tablet formulations of the active ingredient (equivalent to 5 mg anhydrous free 4-amino-1-hydroxy- butylidene-1,1-bisphosphonic acid per tablet) were prepared under different conditions with differing excipients. The direct compression tablets were prepared according to the procedure of Example 1 and the wet granulated tablets were prepared according to the procedure below. The tablets were subjected to stability studies under open dish conditions at 40°C/75% relative humidity. The following observations were noted:

2149052

- 14 -

1. Tablet discoloration occurred within 2 weeks in formulations which were manufactured by wet granulation and contained anhydrous lactose.

5 2. Tablet discoloration occurred within 4 weeks in formulations which were manufactured by wet granulation and contained hydrous lactose.

3. There was no tablet discoloration after 4 weeks in formulations which manufactured as a direct compression (dry mix) formulation. Assay of the active ingredient confirmed that there was no  
10 loss of potency or formation of degradates over the same time period. Table I demonstrates the stability characteristics of the direct compression formulation as compared to a wet granulated formulation.

15 TABLE I

Three-Month Data of 5 mg Probe Stability Lots

(a). Direct Compression

20	<u>Condition</u>	<u>Assay, % initial</u>			
		<u>Open Dish</u>		<u>HDPE/CRC Bottle</u>	
		<u>cpd</u>	<u>Adduct</u>	<u>MK-0217</u>	<u>Adduct</u>
25	40°C	101.2%	----	99.8%	----
	40°C/75% RH	102.9%	----	98.5%	----
				100.9%*	----
	60°C	100.6%	----	101.3%	----
	RT/90% RH	103.5%	----	102.1%	----

30 \* Duplicate value

- 15 -

## (b). Wet Granulation

	<u>Condition</u>	<u>Assay, % initial</u>			
		<u>Open Dish</u>		<u>HDPE/CRC Bottle</u>	
		<u>epd</u>	<u>Adduct</u>	<u>MK-0217</u>	<u>Adduct</u>
5	40°C	99.7%	----	97.0%	----
				100.0%**	----
10	40°C/75% RH	84.1%	15.9%	94.6%	5.6%
				99.7%**	----
	60°C	92.6%	trace***	94.3%	trace***
				94.6%**	trace
	RT/90% RH	101.4%	----	100.4%	----
15				99.4%	----

\*\* With desiccant

\*\*\* Trace indicates that the adduct peak is detectable (~5%) but not quantifiable under experimental conditions.

cpd = 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid

20

Process for wet granulation tablets

1) Anhydrous lactose, alendronate sodium and microcrystalline cellulose was mixed in a suitable size blender.

25

2) The blend was strained through mesh #30 followed by remixing for an additional time.

3) The powder mixture was granulated with an adequate quantity of water until caking occurred.

30

4) The wet mass was poured through a screen #5.

2149052

- 16 -

5) The wet sized granulation was dried in a forced air dryer at 40-50°C until the loss on drying was less than 2% at 105°C.

5 6) The dry granulation was sized through a suitable screen.

7) The dry sized granulation was mixed with croscarmellose sodium followed by magnesium stearate.

10 8) Tablets were compressed using the labeled granulation.

15 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the casual variations, adaptations, modifications, deletions, or additions of procedures and protocols described herein, as come within the scope of the following claims and its equivalents.

20

25

30